

## AN ENERGY DISPERSIVE X-RAY FLUORESCENCE STUDY OF SOME NEAR EASTERN OBSIDIANS

Mark E. HALL, Dr.\* and Steven SHACKLEY, Dr.\*\*

### Introduction

For over twenty five years the chemical sourcing of obsidian to study patterns of trade and exchange has been an important aspect of Near Eastern archaeology [Benedict et. al. 1980; Blackman 1984; Cann and Renfrew 1964; Cauvin et. al. 1986; Francaviglia 1990; Perlman and Yellin 1980; Renfrew, Dixon and Cann 1966, 1968; Wright 1969; Wright and Gordus 1969; Zarins 1990]. In a majority of these studies either neutron activation analysis or some form of atomic or optical spectroscopy was used to obtain the chemical composition of the obsidian. Both of these analytical methods can be costly and time consuming; also, in the case of spectroscopy, an actual sample of the material needs to be taken from the artifact.

One purpose of this paper is to illustrate the utility of energy dispersive x-ray fluorescence (EDXRF) in the characterization of obsidian from the Near East. This non-destructive method of analysis is low-cost, can be run quickly, and yields quantitative data. It should be stressed that EDXRF is not a new analytical technique, but one used extensively by archaeologists and geologists studying obsidian particularly in the Americas [Bouey 1991; Fowler et. al. 1989; Hughes 1984, 1988; Hughes and Smith 1993; Jack and Carmichael 1969; Shackley 1988, 1991, 1992].

Another purpose of this paper is to present some new chemical analyses of obsidian artifacts from three Near Eastern sites. The analyses performed on surface finds from Hamoukar and Hirbet Tueris in northeastern Syria and Umm Dabighiyah in Iraq (Figure 1).

### X-Ray Fluorescence Spectrometry

The technique of x-ray fluorescence involves bombarding a material with an x-ray beam of known energy and wavelength. The primary x-ray beam displaces the electrons in the K, L, and M orbitals of the target atoms. The displaced electrons are then replaced by electrons from the outer orbitals of the target atoms; secondary or fluorescent x-rays are released as the outer orbital electrons loose energy to fill the inner orbitals. Each element in a material has a distinct secondary x-ray spectra. For the more interested reader, Williams [1987: 16–46] provides an in-depth overview of the physics of x-ray fluorescence.

After the secondary x-rays are produced, they can be collected and analyzed using one of two elemental methods. Energy dispersive x-ray fluorescence (EDXRF) machines utilize silicon based semi-conductors to detect and measure the intensity of the fluorescent x-rays [Hampel 1984: 21; Potts 1987: 286–303; Williams 1987: 110–120]. Wavelength dispersive x-ray fluorescence (WDXRF) machines utilize a crystal to diffract and separate the secondary x-rays to an x-ray detector. A comparison of the two methods are reviewed in Potts [1987: 299, 300] and Williams [1987: 100–120].

EDXRF is particularly well-suited for obsidian studies since it can accurately measure elements with atomic numbers 11 through 41 and some of the rare earth elements [Hampel 1984: 21, 22; Potts 1987: 312, 313]. The detection limit of EDXRF for niobium (Nb), rubidium (Rb), strontium (Sr), yttrium (Y), and zirconium (Zr), all trace elements used in sourcing obsidian (see below), is as low as 3.5 ppm to 6.0 ppm.

---

\* Archaeological Research Facility, Kroeber Hall, University of California, Berkeley, CA 94720–3710

\*\* Phoebe Hearst Museum of Anthropology, Kroeber Hall, University of California, Berkeley, CA 94720–3712




Fig. 1 Map of the sites and obsidian sources mentioned in the text

Methods of Analysis

The trace elements used to source obsidian in EDXRF and NAA studies are known as “incompatible elements” since they are incompatible with the solid-phase constituents in high temperature silicic melts and are most stable in the resulting glass phase [Cann 1983; Ferrara and Treuil 1975; Hess 1991; Mahood and Stimac 1990; Zielinski et. al. 1977]. They generally do not covary within a single silicic melt and their concentrations are unique for each volcanic event. The incompatible elements include niobium (Nb), rubidium (Rb), strontium (Sr), titanium (Ti), yttrium (Y), and zirconium (Zr). Other major and trace elements, such as chromium (Cr), cobalt (Co), iron (Fe), gallium (Ga), nickel (Ni), and sodium (Na), can vary extensively in a single obsidian source since they are easily absorbed into the solid phase within a silicic melt. Furthermore, iron and sodium can be lost due to hydration and secondary crystallization of the obsidian [MacDonald and Bailey 1973: N2].

The trace element analyses were performed in the Department of Geology and Geophysics, University of California, Berkeley, using a Spectrace™ 440 [United Scientific Corporation] energy dispersive x-ray fluorescence spectrometer. The spectrometer is equipped with a rhodium x-ray tube, a 50 kV x-ray generator, and a Tracor X-ray (Spectrace™) TX 6100 x-ray analyzer using an IBM PC based micro-processor and Tracor reduction software. The x-ray tube was operated at 30 kV, 0.20 mA, using a .127 mm rhodium primary beam filter in an air path at 250 seconds livetime to generate x-ray intensity data. Secondary x-ray intensities were converted to concentration values by employing a least-squares calibration line established for each element from the analysis of up to 26 international rock standards certified by the U.S. Bureau of Standards, the U.S. Geological Survey, Canadian Centre for Mineral and Energy

Technology, the Centre de Recherches Pétrographiques et Géochimiques in France, and the Geological Survey of Japan.

To insure machine calibration, a known standard was run with the obsidian samples. Table IA shows a comparison between values recommended for RGM-1, a U.S. Geological Survey rhyolite rock standard, and the results obtained for it from this study. This data indicates that the machine accuracy is quite high. The maximum amount of error in each elemental reading is listed in Table IB.

Artifact Analyses

The obsidian artifacts in this study came from Hamoukar (Hamukar) and Hirbet Tueris in northeastern Syria and Umm Dabighiyah in Iraq [see Figure 1]. Both Hamoukar and Hirbet Tueris are unexcavated tells in the Habur River basin east of the al-Radd marsh and close to the border with Iraq [Dobel 1978: 103, 108, 194–197; van Liere 1963]. Hamukar is believed by van Liere [1963: 114–120] to be the capital of the Mitanni state, Wassukanni. Hirbet Tueris is a smaller tell near Hamukar. From surface finds of pottery, occupation continued at Hirbet Tueris through the second millennium BC [Dobel 1978: 103]. Umm Dabighiyah is a small tell located in the dry steppe zone of the Jazira. Excavations were conducted there in the early 1970s [Kirkbride 1972, 1973a, 1973b]. The finds indicate that the site was occupied during the sixth millennium BC.

The artifacts collected from Hamukar (Nos. 14511–14518, 14520, 14521) and Hirbet Tueris (Nos. 14105–14115) are retouched blades ranging in length from 1.5 cm to 6.0 cm. Blades 14511 and 14514 both contain cortex on one side. The artifacts from Umm Dabighiyah consist of two small re-touched flakes and a piece of debitage.

Table II contains the concentration (in ppm) of the incompatible elements analyzed by EDXRF for this study. Table III contains the concentration of the minor and trace elements in the artifacts. To see how many obsidian sources are present, selected incompatible elements are plotted against each other in Figures 2 through 5. Figure 2 is a plot of Sr versus Zr, Figure 3 is a plot of Y versus Zr, and Figure 4 is a plot of Nb versus Rb, and Figure 5 is a plot of Sr versus Y.

In all four graphs it is clear that the obsidian used at these sites came from three distinct sources.

Table IA

| Sample                   | Ti                | Mn              | Fe                 | Rb             | Sr             | Y            | Zr            | Nb           |
|--------------------------|-------------------|-----------------|--------------------|----------------|----------------|--------------|---------------|--------------|
| RGM-1 (Govindaraju 1989) | 1600              | 279             | 12998              | 149            | 108            | 25           | 219           | 8.9          |
| RGM-1 (this study)       | 1514.29<br>+/-108 | 234.72<br>+/-28 | 13631.28<br>+/-457 | 147.43<br>+/-2 | 106.14<br>+/-4 | 23.5<br>+/-2 | 227.2<br>+/-5 | 8.87<br>+/-5 |

X-ray fluorescence concentrations for selected trace elements of RGM-1. The +/- values represent the first standard deviation computations for the group of measurements. All values are in parts per million (ppm) as reported in Govindaraju (1989) and this study. RGM-1 is a U.S. Geological Survey rhyolite (obsidian) rock standard.

Table IB

| Element | Error (+/-) | Element | Error (+/-) |
|---------|-------------|---------|-------------|
| Nb      | 5.0 ppm     | Fe      | .05%        |
| Rb      | 3.5 ppm     | Mn      | 35 ppm      |
| Sr      | 5.5 ppm     | Ti      | 118 ppm     |
| Y       | 3.0 ppm     | Zn      | 7 ppm       |
| Zr      | 7.5 ppm     |         |             |

Listed above are the maximum amount of measurement errors for each elemental concentration listed in Table II and Table III.

Table II

| Sample number | Nb   | Rb    | Sr   | Th   | Y     | Zr     |
|---------------|------|-------|------|------|-------|--------|
| 14511         | 53.8 | 220.9 | 7.5  | 17.6 | 119.4 | 1185.1 |
| 14512         | 34.8 | 200.5 | 21.7 | 23.6 | 50.6  | 266.3  |
| 14513         | 68.3 | 241.5 | 7.3  | 31.3 | 129.9 | 1263.0 |
| 14514         | 65.2 | 239.8 | 7.2  | 32.8 | 134.9 | 1254.1 |
| 14515         | 63.7 | 225.7 | 4.9  | 27.6 | 123.2 | 1193.6 |
| 14516         | 58.0 | 222.8 | 6.7  | 29.6 | 119.1 | 1148.3 |
| 14517         | 61.7 | 238.3 | 5.3  | 29.7 | 127.7 | 1253.3 |
| 14518         | 62.0 | 218.9 | 7.0  | 25.8 | 124.0 | 1180.9 |
| 14520         | 65.9 | 236.0 | 4.4  | 27.9 | 122.6 | 1240.2 |
| 14521         | 64.3 | 232.5 | 5.0  | 25.8 | 124.0 | 1226.5 |
| 145105        | 60.0 | 221.8 | 13.5 | 32.9 | 124.8 | 1190.5 |
| 145106        | 58.2 | 229.3 | 6.4  | 29.1 | 127.2 | 1208.5 |
| 145107        | 63.6 | 219.7 | 5.5  | 25.8 | 125.1 | 1179.4 |
| 145108        | 58.9 | 212.0 | 8.9  | 31.4 | 117.5 | 1155.3 |
| 145109        | 63.9 | 230.7 | 12.8 | 30.2 | 123.6 | 1204.2 |
| 145110        | 60.6 | 232.4 | 9.0  | 32.9 | 125.3 | 1214.9 |
| 145111        | 57.0 | 228.1 | 8.9  | 31.1 | 121.2 | 1165.3 |
| 145112        | 56.9 | 229.8 | 7.2  | 23.4 | 123.6 | 1206.8 |
| 145113        | 63.2 | 229.4 | 5.5  | 26.8 | 126.2 | 1226.6 |
| 145114        | 65.4 | 232.0 | 6.0  | 32.4 | 127.0 | 1223.5 |
| 145115        | 62.8 | 235.5 | 6.3  | 27.8 | 124.7 | 1228.6 |
| UD1           | 55.6 | 220.3 | 4.1  | 40.0 | 124.5 | 1122.7 |
| UD2           | 13.9 | 223.2 | 51.7 | 31.6 | 30.1  | 297.7  |
| UD3           | 62.4 | 204.5 | 5.4  | 37.2 | 122.5 | 1086.3 |

Concentration of incompatible and trace elements in the artifacts from Hamakar (Nos. 14511 to 14518, 14520, 14521), Ḣirbet Ṫueris (Nos. 14105 to 14115), and Umm Dabighiyah (Nos. UD1 to UD3). All concentrations are listed in parts per million (ppm).

Table III

| Sample number | Fe (%) | Mn    | Ti     | Zn    |
|---------------|--------|-------|--------|-------|
| 14511         | 2.08   | 393.9 | 900.4  | 174.8 |
| 14512         | 1.18   | 466.8 | 652.0  | 92.4  |
| 14513         | 2.49   | 526.7 | 993.1  | 192.4 |
| 14514         | 2.60   | 511.3 | 1126.2 | 197.5 |
| 14515         | 2.20   | 458.7 | 950.8  | 175.7 |
| 14516         | 2.34   | 429.3 | 929.6  | 201.2 |
| 14517         | 2.44   | 491.3 | 1039.1 | 193.7 |
| 14518         | 2.28   | 434.4 | 941.4  | 183.1 |
| 14520         | 2.39   | 484.1 | 991.9  | 188.7 |
| 14521         | 2.37   | 525.8 | 1001.9 | 186.9 |
| 145105        | 2.14   | 421.2 | 991.5  | 169.6 |
| 145106        | 2.50   | 492.0 | 960.4  | 202.2 |
| 145107        | 2.23   | 413.1 | 918.5  | 186.6 |
| 145108        | 2.15   | 413.5 | 906.7  | 190.2 |
| 145109        | 2.24   | 450.2 | 967.8  | 179.5 |
| 145110        | 2.32   | 451.1 | 1013.4 | 181.2 |
| 145111        | 2.36   | 464.9 | 1108.6 | 209.5 |
| 145112        | 2.34   | 452.8 | 994.8  | 202.8 |
| 145113        | 2.37   | 462.7 | 1064.2 | 190.5 |
| 145114        | 2.30   | 524.3 | 1062.9 | 193.0 |
| 145115        | 2.30   | 459.4 | 883.8  | 179.4 |
| UD1           | 3.38   | 572.7 | 2065.9 | 220.1 |
| UD2           | 1.53   | 266.7 | 1359.4 | 65.7  |
| UD3           | 2.02   | 334.6 | 973.0  | 174.1 |

Concentration of minor and trace elements in the artifacts from Hamakar (Nos. 14511 to 14518, 14520, 14521), Ḣirbet Ṫueris (Nos. 14105 to 14115), and Umm Dabighiyah (Nos. UD1 to UD3). All concentrations, except iron (Fe) are listed in parts per million (ppm). The iron concentration is in %.




Fig. 2 Plot of Sr concentration versus Zr concentration




Fig. 3 Plot of Y concentration versus Zr concentration




Fig. 4 Plot of Nb concentration versus Rb concentration




Fig. 5 Plot of Y concentration versus Sr concentration

Twenty-two out of 24 artifacts came from a single source; the remaining two artifacts each came from a distinct source.

These sources can be identified by using the published data in Benedict et. al. [1980], Blackman [1984: 42–49] and Cauvin et. al. [1986: 92]. On the basis of the iron, rubidium, thorium, and zinc concentrations, the majority of the artifacts are made from obsidian coming from one of the Nemrut Dag sources. Due to the lack of data on two of the obsidian flows at Nemrut Dag, it cannot be ascertained which of the four sources provided the obsidian. On the basis of the iron, manganese, strontium, titanium, yttrium, zinc and zirconium concentrations, sample UD2, the debitage from Umm Dabighiyah, is made from obsidian coming from source B at Bingöl, Turkey. Blade 14512 from Hamukar was made of obsidian from an unknown source. Its composition is similar to obsidian artifacts found at Zarnaki Tepe north of Lake Van [Blackman 1984: 48–49].

### Discussion

As can be expected, the obsidian used by these three sites came from eastern Turkey. Past research [Blackman 1984: 34–36; Renfrew, Dixon and Cann 1966: 40–49] has shown that most of the obsidian used in Mesopotamia and the Iranian highlands during the 8th millennium BC through the mid-3rd millennium BC came from Nemrut Dag. The Bingöl sources and the source supplying Zarnaki Tepe were subsidiary to Nemrut Dag at this time. After the 3rd millennium BC, Nemrut Dag lost its prominence in the Iranian highlands and other sources began to be used there. It is uncertain what obsidian sources were in use in Mesopotamia from the mid-3rd millennium BC onwards.

Previous sourcing work on the obsidian from Umm Dabighiyah used color only [Kirkbride 1972: 11; Mellart 1975: 138]. On the basis of color, it was determined that the obsidian came from the Lake Van area. The EDXRF analyses provide more specific information, indicating that the obsidian came from a source at Nemrut Dag and Bingöl source B.

The analyses of the obsidian from Hamoukar and Hirbet Tueris suggest that Nemrut Dag could have been a major source of obsidian as late as the 3rd and 2nd millennium BC for the inhabitants of the Habur River basin. Lake Van was just outside the boundary of the Mitanni province of Alshe [Roaf 1990: 134]. It is quite possible that the Mitanni had ready access to the obsidian at Nemrut Dag, but more research is needed on this issue.

The presence of cortex on two of the blades from Hamoukar and Hirbet Tueris and the debitage from Umm Dabighiyah needs to be noted. These two features suggest that the obsidian nodules were traded rather than the finished tools.

### Conclusion

Over 90% of the obsidian artifacts analyzed were made from obsidian procured from Nemrut Dag. The remainder of the artifacts were made of obsidian from Bingöl and an unknown source.

Even though obsidian sourcing studies have been undertaken for nearly thirty years, research still remains to be done in three major areas. First, the unknown obsidian sources need to be located and characterized. More intensive chemical studies need to be done on obsidian from Hotmis Dag, Lake Sevan, Nemrut Dag, and Suphan Dag. Finally, there needs to be more analyses of obsidian from 3rd and 2nd millennium sites in Mesopotamia. In all of these areas EDXRF can be an effective analytical tool in the investigation of Near Eastern archaeological obsidian.

### Acknowledgements

The Phoebe Hearst Museum of Anthropology (103 Kroeber Hall, UCB, Berkeley, CA 94720–3712) is

acknowledged for their assistance and cooperation with this project. The majority of the artifacts analyzed in this study are part of the museum's Near Eastern collection.

### Bibliography

- Benedict, Peter, Adon A. Gordus, Mehmet Özdoğan, and Gary Wright.  
 1980 "Location and Chemical Identification of Some Obsidian Sources in the Aksaray-Nevşehir-Nigde Region, Central Anatolia," *Prehistoric Research in Southeastern Anatolia* by Halet Çambel and Robert J. Braidwood, Edebiyat Fakültesi Basımevi, İstanbul, pp. 239–256.
- Blackman, M. James.  
 1984 "Provenance Studies of Middle Eastern Obsidian from Sites in Highland Iran," *Archaeological Chemistry III* edited by Joseph B. Lambert, Advances in Chemistry 205, American Chemical Society, Washington, D.C., pp. 19–50.
- Bouey, Paul D.  
 1991 "Recognizing the Limits of Archaeological Applications of Non-Destructive Energy-Dispersive X-Ray Fluorescence Analysis of Obsidians," *Materials Issues in Art and Archaeology II* edited by Pamela B. Vandiver, James Druzik, and George Seagan Wheeler, Materials Research Society Symposium Proceedings Vol. 185, Materials Research Society, Pittsburgh, pp. 301–320.
- Cauvin, M.-C., N. Balkan, Y. Besnus, and F. Saroglu.  
 1986 "Origine de L'obsidienne de Cafer Hoyuk [Turquie]: Premiers Résultats," *Paléorient*, Vol. 12, No. 2, pp. 89–97.
- Cann, J.R.  
 1983 "Petrology of Obsidian Artefacts," *The Petrology of Archaeological Artefacts* edited by D.R.C. Kempe and A.P. Harvey, Oxford University Press, Oxford, pp. 227–255.
- Cann, J.R. and Colin Renfrew.  
 1964 "The Characterization of Obsidian and its application to the Mediterranean Region," *Proceedings of the Prehistoric Society*, Vol. 30, pp. 111–133.
- Dixon, J.E.  
 1976 "Obsidian Characterization Studies in the Mediterranean and Near East," *Advances in Obsidian Glass Studies* edited by R.E. Taylor, Noyes Press, Park Ridge, New Jersey, pp. 288–333.
- Dobel, Allan George.  
 1978 *The Location of Wassukanni: An Analysis of Archaeological and Textual Source Materials*, unpublished Ph. D. dissertation, University of California, Berkeley, 1978.
- Ferrara, G. and M. Treuil.  
 1975 "Petrological Implications of Trace Element and Sr Isotope Distributions in Basalt-Pantellerite Series," *Bulletin Volcanologique*, Volume 38, No. 3, pp. 548–574.
- Fowler, Jr., William R., Arthur A. Demarest, Helen V. Michel, Frank Asaro, Fred Stross.  
 1989 "Sources of Obsidian from El Mirador, Guatemala: New Evidence on PreClassic Maya Interaction," *American Anthropologist*, Vol. 91, pp. 158–168.
- Francaviglia, Vincenzo.  
 1990 "Obsidian Sources in Ancient Yemen." In *The Bronze Age Culture of Hawlan At-Tiyal and Al-Hada* by Alessandro De Maigret, ISMEO, Rome, pp. 129–136.
- Govindaraju, K.  
 1989 "1989 Compilation of Working Values and Sample Description for 272 Geo-standards," *Geostandards Newsletter*, Vol. 13.
- Hampel, Joachim H.  
 1984 "Technical Considerations in X-Ray Fluorescence Analysis of Obsidians," *Obsidian Studies in the Great Basin* edited by Richard E. Hughes, Contributions of the University of California Archaeological Research Facility Number 45, Archaeological Research Facility, University of California at Berkeley, Berkeley, pp. 21–25.
- Hess, Paul C.  
 1991 "The Role of High Field Strength Cations in Silicate Melts," *Physical Chemistry of Magmas* edited by L.I. Perchuk and I. Kushiro, Advances in Physical Geochemistry Vol. 9, Springer-Verlag, New York, pp. 152–191.
- Hughes, Richard.  
 1984 "Obsidian Sourcing Studies in the Great Basin: Problems and Prospects," *Obsidian Studies in the Great Basin* edited by Richard E. Hughes, Contributions of the University of California Archaeological Research Facility Number 45, Archaeological Research Facility, University of California at Berkeley, pp. 1–19.  
 1988 "The Coso Volcanic Field Reexamined: Implications for Obsidian Sourcing and Hydration Dating Research," *Geoarchaeology*, Vol. 3, No. 4, pp. 253–265.

- Hughes, Richard and Robert L. Smith.
  - 1993 "Archaeology, Geology, and Geochemistry in Obsidian Provenance Studies," *Effects of Scale on Archaeological and Geoscientific Perspectives* edited by J.K. Stein and A.R. Linse, Geological Society of America Special Paper 283, pp. 79–91.
- Jack, R.N. and I.S.E. Carmichael.
  - 1969 "The Chemical Fingerprinting of Acid Volcanic Rocks," *California Division of Mines and Geology, Special Report 100*, pp. 17–32.
- Kirkbride, Diana.
  - 1972 "Umm Dabaghiyah 1971: A Preliminary Report," *Iraq*, Volume 34, pp. 3–15.
  - 1973a "Umm Dabaghiyah 1972: A Second Preliminary Report," *Iraq*, Volume 35, pp. 1–7.
  - 1973b "Umm Dabaghiyah 1973: A Third Preliminary Report," *Iraq*, Volume 35, pp. 205–209.
- MacDonald, Ray and D.K. Bailey.
  - 1973 "The Chemistry of Peralkaline Oversaturated Obsidians," *Data of Geochemistry, Chapter N: Chemistry of Igneous Rocks* edited by Michael Fleischer, Geological Survey Professional Paper 440-N-1, U.S. Government Printing Office, Washington, D.C.
- Mahood, Gail A. and James A. Stimac.
  - 1990 "Trace-element Partitioning in Pantellerites and Trachytes," *Geochimica et Cosmochimica Acta*, Vol. 54, pp. 2257–2276.
- Mellart, James.
  - 1975 *The Neolithic of the Near East*, Charles Scribner's and Sons, New York.
- Perlman, Isadore and Joseph Yellin.
  - 1980 "The Province of Obsidian from Neolithic Sites in Israel," *Israel Exploration Journal*, Vol. 30, Nos. 1–2, pp. 83–88.
- Potts, P.J.
  - 1987 *A Handbook of Silicate Rock Analysis*, Chapman and Hall, New York.
- Renfrew, Colin, J.E. Dixon, and J.R. Cann.
  - 1966 "Obsidian and Early Cultural Contact in the Near East," *Proceedings of the Prehistoric Society*, Vol. 32, pp. 30–72.
  - 1968 "Further Analysis of Near Eastern Obsidians," *Proceedings of the Prehistoric Society*, Vol. 34, pp. 319–331.
- Roaf, Michael.
  - 1990 *Cultural Atlas of Mesopotamia and the Ancient Near East*, Facts on File, Inc., New York.
- Shackley, M. Steven.
  - 1988 "Sources of Archaeological Obsidian in the Southwest: An Archaeological, Petrological, and Geochemical Study," *American Antiquity*, Volume 53, No. 4, pp. 752–772.
  - 1991 "Tank Mountains Obsidian: A Newly Discovered Archaeological Obsidian Source in East-Central Yuma County, Arizona," *Kiva*, Vol. 57, No. 1, pp. 17–25.
  - 1992 "The Upper Gila River Gravels as an Archaeological Obsidian Source Region: Implications for Models of Exchange and Interaction," *Geoarchaeology*, Vol. 7, No. 4, pp. 315–326.
- van Liere, W.J.
  - 1963 "Capitals and Citadels," *Annales Archéologiques de Syrie*, Vol. 13, pp. 109–122.
- Williams, K.L.
  - 1987 *Introduction to X-Ray Spectrometry*. Allen and Unwin, London.
- Wright, Gary A.
  - 1969 *Obsidian Analyses and Prehistoric Near Eastern Trade: 7500 to 3500 B.C.* Anthropological Papers No. 37, University of Michigan, Ann Arbor.
- Wright, Gary A. and A.A. Gordus.
  - 1969 "Source Areas for Obsidian Recovered at Munhata, Beisamoun, Hazorea and El-Khiam," *Israel Exploration Journal*, Vol. 19, No. 2, pp. 79–88.
- Zarins, Juris.
  - 1990 "Obsidian and the Red Sea Trade: Prehistoric Aspects," *South Asian Archaeology 1987* edited by Maurizio Taddei and Pierfrancesco Callieri, ISMEO, Rome, pp. 507–541.
- Zielinski, Robert, Peter Lipman and Hugh Millard.
  - 1977 "Minor-element Abundances in Obsidian, Perlite, and Felsite of Calc-Alkalic Rhyolites," *American Mineralogist*, Vol. 62, pp. 426–437.